


1999 - 2015 Average Death Rates for Prescription* Drug Overdose by State

[View Data Tables](#)

(move cursor over state abbreviation to display death rates by year)



PDMP TTAC
Prescription Drug Monitoring Program Training and Technical Assistance Center

Age adjusted rate per
100,000 population


- <= 4.9 death rate
- 5.0 to 9.9 death rate
- 10.0 to 14.9 death rate
- >= 15.0 death rate
- Data unreliable or suppressed **

*Prescription Drug - refers to a pharmaceutical drug that requires a medical prescription to be dispensed; includes Schedule II - V controlled substances and legend drugs (i.e., amphetamines, barbiturates, opiates, antibiotics, statin drugs). Excludes illicit and unspecified drugs.

**Death rates based on counts of less than twenty (death count < 20) are flagged as "Unreliable". A death rate based on fewer than 20 deaths has a relative standard error (RSE(R)) of 23 percent or more. A RES(R) of 23 percent is considered statistically unreliable. Death counts and death rates are "Suppressed" when the figure represents zero to nine (0-9) persons.

Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Multiple Cause of Death 1999-2015 on CDC WONDER Online Database, released 2016. Data are from the Multiple Cause of Death Files, 1999-2015, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program.

1999 Death Rates for Prescription* Drug Overdose by State





*Prescription Drug - refers to a pharmaceutical drug that requires a medical prescription to be dispensed; includes Schedule II - V controlled substances and legend drugs (i.e., amphetamines, barbiturates, opiates, antibiotics, statin drugs). Excludes illicit and unspecified drugs.

**Death rates based on counts of less than twenty (death count < 20) are flagged as "Unreliable". A death rate based on fewer than 20 deaths has a relative standard error (RSE(R)) of 23 percent or more. A RES(R) of 23 percent is considered statistically unreliable. Death counts and death rates are "Suppressed" when the figure represents zero to nine (0-9) persons.

Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Multiple Cause of Death 1999-2014 on CDC WONDER Online Database, released 2015. Data are from the Multiple Cause of Death Files, 1999-2014, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program.

2000 Death Rates for Prescription* Drug Overdose by State




*Prescription Drug - refers to a pharmaceutical drug that requires a medical prescription to be dispensed; includes Schedule II - V controlled substances and legend drugs (i.e., amphetamines, barbiturates, opiates, antibiotics, statin drugs). Excludes illicit and unspecified drugs.

**Death rates based on counts of less than twenty (death count < 20) are flagged as "Unreliable". A death rate based on fewer than 20 deaths has a relative standard error (RSE(R)) of 23 percent or more. A RES(R) of 23 percent is considered statistically unreliable. Death counts and death rates are "Suppressed" when the figure represents zero to nine (0-9) persons.

Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Multiple Cause of Death 1999-2014 on CDC WONDER Online Database, released 2015. Data are from the Multiple Cause of Death Files, 1999-2014, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program.

2001 Death Rates for Prescription* Drug Overdose by State



Age adjusted rate per
100,000 population


- ≤ 4.9 death rate
- 5.0 to 9.9 death rate
- 10.0 to 14.9 death rate
- ≥ 15.0 death rate
- Data unreliable or suppressed **

*Prescription Drug - refers to a pharmaceutical drug that requires a medical prescription to be dispensed; includes Schedule II - V controlled substances and legend drugs (i.e., amphetamines, barbiturates, opiates, antibiotics, statin drugs). Excludes illicit and unspecified drugs.

**Death rates based on counts of less than twenty (death count < 20) are flagged as "Unreliable". A death rate based on fewer than 20 deaths has a relative standard error (RSE(R)) of 23 percent or more. A RES(R) of 23 percent is considered statistically unreliable. Death counts and death rates are "Suppressed" when the figure represents zero to nine (0-9) persons.

Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Multiple Cause of Death 1999-2014 on CDC WONDER Online Database, released 2015. Data are from the Multiple Cause of Death Files, 1999-2014, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program.

2002 Death Rates for Prescription* Drug Overdose by State



Age adjusted rate per 100,000 population


- <= 4.9 death rate
- 5.0 to 9.9 death rate
- 10.0 to 14.9 death rate
- >= 15.0 death rate
- Data unreliable or suppressed **

*Prescription Drug - refers to a pharmaceutical drug that requires a medical prescription to be dispensed; includes Schedule II - V controlled substances and legend drugs (i.e., amphetamines, barbiturates, opiates, antibiotics, statin drugs). Excludes illicit and unspecified drugs.

**Death rates based on counts of less than twenty (death count < 20) are flagged as "Unreliable". A death rate based on fewer than 20 deaths has a relative standard error (RSE(R)) of 23 percent or more. A RES(R) of 23 percent is considered statistically unreliable. Death counts and death rates are "Suppressed" when the figure represents zero to nine (0-9) persons.

Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Multiple Cause of Death 1999-2014 on CDC WONDER Online Database, released 2015. Data are from the Multiple Cause of Death Files, 1999-2014, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program.

2003 Death Rates for Prescription* Drug Overdose by State



Age adjusted rate per 100,000 population


- <= 4.9 death rate
- 5.0 to 9.9 death rate
- 10.0 to 14.9 death rate
- >= 15.0 death rate
- Data unreliable or suppressed **

*Prescription Drug - refers to a pharmaceutical drug that requires a medical prescription to be dispensed; includes Schedule II - V controlled substances and legend drugs (i.e., amphetamines, barbiturates, opiates, antibiotics, statin drugs). Excludes illicit and unspecified drugs.

**Death rates based on counts of less than twenty (death count < 20) are flagged as "Unreliable". A death rate based on fewer than 20 deaths has a relative standard error (RSE(R)) of 23 percent or more. A RES(R) of 23 percent is considered statistically unreliable. Death counts and death rates are "Suppressed" when the figure represents zero to nine (0-9) persons.

Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Multiple Cause of Death 1999-2014 on CDC WONDER Online Database, released 2015. Data are from the Multiple Cause of Death Files, 1999-2014, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program.

2004 Death Rates for Prescription* Drug Overdose by State




*Prescription Drug - refers to a pharmaceutical drug that requires a medical prescription to be dispensed; includes Schedule II - V controlled substances and legend drugs (i.e., amphetamines, barbiturates, opiates, antibiotics, statin drugs). Excludes illicit and unspecified drugs.

**Death rates based on counts of less than twenty (death count < 20) are flagged as "Unreliable". A death rate based on fewer than 20 deaths has a relative standard error (RSE(R)) of 23 percent or more. A RES(R) of 23 percent is considered statistically unreliable. Death counts and death rates are "Suppressed" when the figure represents zero to nine (0-9) persons.

Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Multiple Cause of Death 1999-2014 on CDC WONDER Online Database, released 2015. Data are from the Multiple Cause of Death Files, 1999-2014, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program.

2005 Death Rates for Prescription* Drug Overdose by State




*Prescription Drug - refers to a pharmaceutical drug that requires a medical prescription to be dispensed; includes Schedule II - V controlled substances and legend drugs (i.e., amphetamines, barbiturates, opiates, antibiotics, statin drugs). Excludes illicit and unspecified drugs.

**Death rates based on counts of less than twenty (death count < 20) are flagged as "Unreliable". A death rate based on fewer than 20 deaths has a relative standard error (RSE(R)) of 23 percent or more. A RES(R) of 23 percent is considered statistically unreliable. Death counts and death rates are "Suppressed" when the figure represents zero to nine (0-9) persons.

Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Multiple Cause of Death 1999-2014 on CDC WONDER Online Database, released 2015. Data are from the Multiple Cause of Death Files, 1999-2014, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program.

2006 Death Rates for Prescription* Drug Overdose by State




*Prescription Drug - refers to a pharmaceutical drug that requires a medical prescription to be dispensed; includes Schedule II - V controlled substances and legend drugs (i.e., amphetamines, barbiturates, opiates, antibiotics, statin drugs). Excludes illicit and unspecified drugs.

**Death rates based on counts of less than twenty (death count < 20) are flagged as "Unreliable". A death rate based on fewer than 20 deaths has a relative standard error (RSE(R)) of 23 percent or more. A RES(R) of 23 percent is considered statistically unreliable. Death counts and death rates are "Suppressed" when the figure represents zero to nine (0-9) persons.

Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Multiple Cause of Death 1999-2014 on CDC WONDER Online Database, released 2015. Data are from the Multiple Cause of Death Files, 1999-2014, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program.

2007 Death Rates for Prescription* Drug Overdose by State




PDMP TTAC
 Prescription Drug Monitoring Program Training and Technical Assistance Center

*Prescription Drug - refers to a pharmaceutical drug that requires a medical prescription to be dispensed; includes Schedule II - V controlled substances and legend drugs (i.e., amphetamines, barbiturates, opiates, antibiotics, statin drugs). Excludes illicit and unspecified drugs.

**Death rates based on counts of less than twenty (death count < 20) are flagged as "Unreliable". A death rate based on fewer than 20 deaths has a relative standard error (RSE(R)) of 23 percent or more. A RES(R) of 23 percent is considered statistically unreliable. Death counts and death rates are "Suppressed" when the figure represents zero to nine (0-9) persons.

Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Multiple Cause of Death 1999-2014 on CDC WONDER Online Database, released 2015. Data are from the Multiple Cause of Death Files, 1999-2014, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program.

2008 Death Rates for Prescription* Drug Overdose by State




*Prescription Drug - refers to a pharmaceutical drug that requires a medical prescription to be dispensed; includes Schedule II - V controlled substances and legend drugs (i.e., amphetamines, barbiturates, opiates, antibiotics, statin drugs). Excludes illicit and unspecified drugs.

**Death rates based on counts of less than twenty (death count < 20) are flagged as "Unreliable". A death rate based on fewer than 20 deaths has a relative standard error (RSE(R)) of 23 percent or more. A RES(R) of 23 percent is considered statistically unreliable. Death counts and death rates are "Suppressed" when the figure represents zero to nine (0-9) persons.

Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Multiple Cause of Death 1999-2014 on CDC WONDER Online Database, released 2015. Data are from the Multiple Cause of Death Files, 1999-2014, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program.

2009 Death Rates for Prescription* Drug Overdose by State



PDMP TTAC
 Prescription Drug Monitoring Program Training and Technical Assistance Center

Age adjusted rate per 100,000 population


- ≤ 4.9 death rate
- 5.0 to 9.9 death rate
- 10.0 to 14.9 death rate
- ≥ 15.0 death rate
- Data unreliable or suppressed **

*Prescription Drug - refers to a pharmaceutical drug that requires a medical prescription to be dispensed; includes Schedule II - V controlled substances and legend drugs (i.e., amphetamines, barbiturates, opiates, antibiotics, statin drugs). Excludes illicit and unspecified drugs.

**Death rates based on counts of less than twenty (death count < 20) are flagged as "Unreliable". A death rate based on fewer than 20 deaths has a relative standard error (RSE(R)) of 23 percent or more. A RES(R) of 23 percent is considered statistically unreliable. Death counts and death rates are "Suppressed" when the figure represents zero to nine (0-9) persons.

Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Multiple Cause of Death 1999-2014 on CDC WONDER Online Database, released 2015. Data are from the Multiple Cause of Death Files, 1999-2014, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program.

2010 Death Rates for Prescription* Drug Overdose by State




*Prescription Drug - refers to a pharmaceutical drug that requires a medical prescription to be dispensed; includes Schedule II - V controlled substances and legend drugs (i.e., amphetamines, barbiturates, opiates, antibiotics, statin drugs). Excludes illicit and unspecified drugs.

**Death rates based on counts of less than twenty (death count < 20) are flagged as "Unreliable". A death rate based on fewer than 20 deaths has a relative standard error (RSE(R)) of 23 percent or more. A RES(R) of 23 percent is considered statistically unreliable. Death counts and death rates are "Suppressed" when the figure represents zero to nine (0-9) persons.

Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Multiple Cause of Death 1999-2014 on CDC WONDER Online Database, released 2015. Data are from the Multiple Cause of Death Files, 1999-2014, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program.

2011 Death Rates for Prescription* Drug Overdose by State




*Prescription Drug - refers to a pharmaceutical drug that requires a medical prescription to be dispensed; includes Schedule II - V controlled substances and legend drugs (i.e., amphetamines, barbiturates, opiates, antibiotics, statin drugs). Excludes illicit and unspecified drugs.


**Death rates based on counts of less than twenty (death count < 20) are flagged as "Unreliable". A death rate based on fewer than 20 deaths has a relative standard error (RSE(R)) of 23 percent or more. A RES(R) of 23 percent is considered statistically unreliable. Death counts and death rates are "Suppressed" when the figure represents zero to nine (0-9) persons.

Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Multiple Cause of Death 1999-2014 on CDC WONDER Online Database, released 2015. Data are from the Multiple Cause of Death Files, 1999-2014, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program.

2012 Death Rates for Prescription* Drug Overdose by State



PDMP TTAC
Prescription Drug Monitoring Program Training and Technical Assistance Center





*Prescription Drug - refers to a pharmaceutical drug that requires a medical prescription to be dispensed; includes Schedule II - V controlled substances and legend drugs (i.e., amphetamines, barbiturates, opiates, antibiotics, statin drugs). Excludes illicit and unspecified drugs.

**Death rates based on counts of less than twenty (death count < 20) are flagged as "Unreliable". A death rate based on fewer than 20 deaths has a relative standard error (RSE(R)) of 23 percent or more. A RES(R) of 23 percent is considered statistically unreliable. Death counts and death rates are "Suppressed" when the figure represents zero to nine (0-9) persons.

Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Multiple Cause of Death 1999-2014 on CDC WONDER Online Database, released 2015. Data are from the Multiple Cause of Death Files, 1999-2014, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program.

2013 Death Rates for Prescription* Drug Overdose by State




*Prescription Drug - refers to a pharmaceutical drug that requires a medical prescription to be dispensed; includes Schedule II - V controlled substances and legend drugs (i.e., amphetamines, barbiturates, opiates, antibiotics, statin drugs). Excludes illicit and unspecified drugs.

**Death rates based on counts of less than twenty (death count < 20) are flagged as "Unreliable". A death rate based on fewer than 20 deaths has a relative standard error (RSE(R)) of 23 percent or more. A RES(R) of 23 percent is considered statistically unreliable. Death counts and death rates are "Suppressed" when the figure represents zero to nine (0-9) persons.

Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Multiple Cause of Death 1999-2014 on CDC WONDER Online Database, released 2015. Data are from the Multiple Cause of Death Files, 1999-2014, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program.


2014 Death Rates for Prescription* Drug Overdose by State




*Prescription Drug - refers to a pharmaceutical drug that requires a medical prescription to be dispensed; includes Schedule II - V controlled substances and legend drugs (i.e., amphetamines, barbiturates, opiates, antibiotics, statin drugs). Excludes illicit and unspecifed drugs.

**Death rates based on counts of less than twenty (death count < 20) are flagged as "Unreliable". A death rate based on fewer than 20 deaths has a relative standard error (RSE(R)) of 23 percent or more. A RES(R) of 23 percent is considered statistically unreliable. Death counts and death rates are "Suppressed" when the figure represents zero to nine (0-9) persons.

Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Multiple Cause of Death 1999-2014 on CDC WONDER Online Database, released 2015. Data are from the Multiple Cause of Death Files, 1999-2014, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program.



2015 Death Rates for Prescription* Drug Overdose by State




*Prescription Drug - refers to a pharmaceutical drug that requires a medical prescription to be dispensed; includes Schedule II - V controlled substances and legend drugs (i.e., amphetamines, barbiturates, opiates, antibiotics, statin drugs). Excludes illicit and unspecified drugs.

**Death rates based on counts of less than twenty (death count < 20) are flagged as "Unreliable". A death rate based on fewer than 20 deaths has a relative standard error (RSE(R)) of 23 percent or more. A RES(R) of 23 percent is considered statistically unreliable. Death counts and death rates are "Suppressed" when the figure represents zero to nine (0-9) persons.

Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Multiple Cause of Death 1999-2015 on CDC WONDER Online Database, released 2016. Data are from the Multiple Cause of Death Files, 1999-2015, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program.


PDMP TTAC
Prescription Drug Monitoring Program Training and Technical Assistance Center


**Age adjusted rate per
100,000 population**

 <= 4.9 death rate

5.0 to 9.9 death rate

10.0 to 14.9 death rate

 ≥ 15.0 death rate

 Data unreliable or suppressed **